



Haemodynamic response during initiation of non-invasive positive pressure ventilation in COPD patients with acute ventilatory failure

M. CONFALONIERI*, P. GAZZANIGA[†], L. GANDOLA*, S. AIOLFI*, R. DELLA PORTA, A. FRISINGHELLI[‡] AND A. BELLONE[§]

*Divisione di *Pneumologia e di [†]Cardiologia, Ospedale Maggiore di Crema, Crema (CR), Italy
Servizio di [‡]Cardiologia e di [§]Fisiopatologia Respiratoria, Ospedale di Passirana di Rho, Rho (MI), Italy*

The aim of this study was to check non-invasively the acute haemodynamic effects of non-invasive positive pressure ventilation (NPPV) initiation in patients with chronic obstructive pulmonary disease (COPD) and acute ventilatory failure (AVF). Nineteen consecutive COPD patients with AVF were evaluated clinically and echocardiographically during spontaneous breathing with O₂ supplementation and during NPPV plus O₂. NPPV was administered with a scheduled inspiratory pressure of 15 cmH₂O and an expiratory pressure of 4 cmH₂O, via facial mask. Arterial blood gas improved significantly (pH and PaCO₂; $P < 0.001$) during NPPV administration in all patients; none had hypotension or acute arrhythmia. Doppler echocardiographic evaluation was feasible in most of the patients (16/18). With reference to baseline values, no significant changes in pulmonary artery pressures and cardiac output (CO) were observed by Doppler echocardiography in most patients. Only four patients (21%) showed a significant reduction ($>15\%$) of CO during NPPV. No correlation was found between decreased CO and baseline data, but three patients showing CO reduction had poor tolerance to mask ventilation and did not improve respiratory rate during NPPV. It was concluded that the initiation of NPPV by facial mask does not alter haemodynamics acutely in most COPD patients with AVF, but individual patients may experience reduction in CO in spite of adequate oxygen saturation levels. This suggests that caution should be used when applying pre-determined and fixed pressures during NPPV. Monitoring haemodynamics by Doppler echocardiography may be useful for early detection of haemodynamic alterations due to NPPV application in patients with AVF.

RESPIR. MED. (1998) 92, 331–337

Introduction

Recent, prospective, randomized trials (1–3) have clearly established non-invasive positive pressure ventilation (NPPV) by nasal or facial mask as a first-line treatment in selected patients with acute ventilatory failure (AVF) due to exacerbation of chronic obstructive pulmonary disease (COPD). In the last few years, a number of controlled and uncontrolled studies, involving more than 500 patients, have appeared in the literature (4), demonstrating very low complication rates associated with NPPV. Moreover, no report exists to date on NPPV-related haemodynamic complications in COPD patients with AVF in the literature. Conversely, it is well known that invasive mechanical ventilation may seriously alter haemodynamic status by decreasing cardiac output and modifying blood flow distribution (5).

The intubation and initiation of positive pressure ventilation in acute exacerbation of COPD (6) is one of the clinical settings which is most likely to trigger haemodynamic compromise. In theory, relatively low support pressures, together with the absence of neuromuscular blockade and sedation, should produce a lower impact of NPPV on haemodynamics. The aim of this study is to assess whether or not the initiation of NPPV by face mask with scheduled inspiratory and expiratory positive pressures can acutely modify haemodynamic status in COPD patients with AVF. For this purpose, we used colour Doppler echocardiography, a non-invasive technique suitable for haemodynamics evaluation in ventilated and COPD patients (7–9).

Methods

SUBJECTS

We studied 19 COPD patients (14 males and five females, mean age 67 ± 9 years) who were admitted for management

Received 1 March 1997 and accepted in revised form 13 November 1997.

Correspondence should be addressed to: M. Confalonieri, via Crotti 24, I-29100 Piacenza, Italy.

of decompensated AVF. COPD diagnosis was confirmed by patient history and physical examination as well as by previous pulmonary function tests. Patients were considered eligible for the study if they conformed with the following criteria:

- respiratory acidosis and AVF (at least two of the following blood gas values: pH <7.30, PaCO₂ >7.90 kPa, PaO₂ <7.98 kPa) due to exacerbation of COPD;
- absence of coma;
- absence of haemodynamic instability (uncontrolled cardiac arrhythmias and/or systolic blood pressure <90 mmHg).

An illness-severity score, simplified acute physiologic score (SAPS) (10), and a neurological score, specifically designed for respiratory patients by Kelly and Matthay (11), were determined for each patient at study entry.

PROTOCOL

At hospital admission, each patient received oxygen supplementation and aerosolized salbutamol (1.25 mg) and ipatropium bromide (0.25 mg) plus beclomethasone (40 mg) in an emergency room setting. After 15–20 min, patients were transferred to the respiratory intermediate intensive care unit (RIICU), for NPPV plus O₂ therapy. Patients received O₂ therapy alone for almost 30 min altogether. NPPV was begun with O₂ supplementation adjusted to give an arterial O₂ saturation of 90–92%. Patients were sequentially evaluated clinically (including non-invasive blood pressure) and echocardiographically during O₂ supplementation at RIICU admission (after 30 min of oxygen supplementation) and during NPPV plus O₂ after a 30 min lapse. Randomization of treatment order was not performed since (a) emergency room, acute respiratory distress patients routinely receive oxygen supplementation before a medical visit, (b) no acute haemodynamic changes in exacerbated COPD and acute respiratory failure patients treated with oxygen supplementation have been reported to date (12–14) and lastly (c) we wished to avoid the hysteric effect caused by positive pressure ventilation (15). Particular care in patient timing management was performed to guarantee from the lack of random order in NPPV application (16). Also, echocardiograph personnel were not aware of the background or aim of the study at the time of Doppler measurements. Furthermore, ethical concerns prohibited baseline evaluation of patients breathing without oxygen supplementation and the inclusion of a non-ventilated control group. Informed consent was given by the patients or next of kin. The study was approved by the Local Ethics Committee.

NON-INVASIVE VENTILATION

NPPV was delivered by a bi-level positive airway pressure (BiPAP® S/T-D, Respiration Inc, Murrissville, PA, U.S.A.) through a facial mask (Gibeck Respiration AB, Upplands-Väsby, Sweden). BiPAP (17) is a flow-triggered device

working as a pressure support (PSV) which can apply external positive end-expiratory pressure (PEEP). We applied BiPAP in spontaneous mode with a scheduled inspiratory pressure support of 15 cmH₂O and an expiratory positive pressure of 4 cmH₂O. We chose these pressure levels based on our clinical experiences of NPPV efficacy (18,19) and on the reported experiences of other authors (20–23). A pneumologist assisted to facilitate patient co-operation at all times. The mask cushion was inflated with about 20 ml of air for perfect adherence to the patient's face and particular care was taken to avoid possible air leaks. Before and during NPPV application, oxygen was administered at the lowest flow to provide an arterial oxygen saturation of 90–92% under oxymetric control (Pulsox 7, Minolta, Avl AG, Switzerland). Patient compliance with ventilation was evaluated by the physician in charge.

ECHOCARDIOGRAPHIC MEASUREMENTS

Two-dimensional, continuous wave (CW), pulsed-wave and colour Doppler echocardiography (Hewlett-Packard 77020A, ultrasound imaging system, Andover, MA, U.S.A.) was performed using 2.5 MHz imaging transducers. Doppler recordings of tricuspid regurgitation velocity and pulmonary flow were attempted from parasternal, apical and subcostal approaches. The 'contrast-enhancement' technique (agitated isotonic saline i.v. injections) was used in selected patients to increase the quality of CW Doppler tracings (24). Maximum velocity of tricuspid regurgitation jet (TR) was measured by CW Doppler imaging. The simplified Bernoulli equation was used to calculate the systolic transtricuspid gradient which was added to mean right atrial pressure (clinically estimated) to predict pulmonary artery systolic pressure (PASP), according to a validated technique (25). The pulsed Doppler technique was used to study pulmonary artery systolic flow velocity, as described by Kitabatake *et al.* (26). Acceleration time (AcT) was defined as the interval between the onset of ejection and peak flow velocity. Right ventricular ejection fraction time (RVET) was defined as the interval between ejection onset to zero flow velocity. We also used the regression equation proposed by Mahan *et al.* (27), which involves AcT, to predict pulmonary artery mean pressure (PAMP).

Blood flow velocity through the aortic valve was assessed by pulsed Doppler echocardiography using a suprasternal approach. Cardiac output (CO) was calculated as the product of the cross-sectional area of the aorta, aortic flow velocity integral and heart rate. Cardiac index (CI) was derived by dividing CO by body surface area. This standard method was extensively validated and used also in COPD patients (28–30). Two-dimensional echocardiographic data, as such the end-diastolic right ventricular:left ventricular dimensions ratio (RVEDD:LVEDD), even though they are useful in a clinical setting (31), were not considered because they are not validated and difficult to obtain in hyperinflated patients.

Since the spontaneous variability of pulmonary artery pressure in COPD patients at rest has a coefficient of

TABLE 1. Patients' characteristics at hospital admission

Patient no.	Age (years)	Sex	Mean BP (mmHg)	PaO ₂ (kPa)	PaCO ₂ (kPa)	pH	Oedema of the legs	RR (breaths min ⁻¹)	HR (beats min ⁻¹)	Kelly score*	SAPS	Baseline† FEV ₁ /FVC (%)
1	72	M	133	5.18	11.7	7.28	=	28	88	2	13	40
2	64	M	106	7.18	7.84	7.35	=	26	96	1	6	31
3	73	M	83	4.78	10.26	7.29	+++	32	102	3	18	48
4	64	M	97	6.46	9.56	7.34	+	28	75	1	8	38
5	51	F	102	5.98	11.71	7.17	=	30	118	2	12	36
6	78	M	110	4.65	10.1	7.19	++	30	120	3	16	?
7	59	M	170	4.89	10.2	7.28	+++	38	90	2	17	29
8	75	M	111	7.71	9.5	7.17	?	36	120	2	14	35
9	58	M	100	4.65	13.73	6.99	+	40	122	3	13	37
10	73	M	103	4.42	13.75	7.15	=	48	99	2	11	36
11	78	M	113	6.38	11.67	7.22	=	34	110	2	15	33
12	72	M	123	6.38	9.84	7.31	+++	32	115	2	17	30
13	67	F	150	4.78	7.71	7.25	=	32	108	1	11	46
14	85	F	93	5.85	11.17	7.19	=	12	130	3	10	?
15	72	F	93	5.18	9.94	7.32	+	28	100	3	9	45
16	56	M	106	6.25	9.27	7.27	=	27	96	1	4	34
17	72	M	93	6.11	9.94	7.32	++	30	100	1	5	40
18	65	F	93	5.18	9.84	7.29	=	30	90	2	8	33
19	54	M	85	6.51	8.15	7.33	=	22	100	1	6	38
Mean	67.7	=	108.6	5.7	10.25	7.25	=	30.6	100.1	2	11.2	37
SD	9.14	=	22.11	0.9	1.76	0.09	=	7.3	14.09	0.8	4.3	5.49
SE	2.1	=	5.07	0.2	0.38	0.02	=	1.6	3.23	0.2	0.9	1.33

*Neurological score by Kelly and Matthay to assess neurological status in patients with acute respiratory failure (11).

†The latest available data before hospital admission.

variation up to 10–6% (32) and since cardiac output measured by echocardiography shows a variability up to 12% (33), we decided that echo Doppler parameter variations of 15% or more would be considered significantly relevant.

REPEATABILITY OF ECHOCARDIOGRAPHIC MEASUREMENTS

Doppler echocardiography was performed by two trained cardiologists from two different institutions. Validation of echocardiograph measurements was previously described by others with regard to CO and PAP (25,30). Repeatability of the technique by the two observers, blinded to each other's results, was previously carried out for eight subjects (four COPD and four healthy subjects; five males; mean age 61±8 years) according to the methodology proposed by Bland and Altman (34).

STATISTICAL ANALYSIS

Individual data and mean ± SD and SE are presented in the tables. Since the data conform with a normal distribution, statistical analysis for values obtained before and during NPPV were performed by paired Student's *t* testing with Bonferonni's adjustment. A linear regression analysis was

used to search for a correlation between haemodynamic and respiratory-clinical data. Statistical significance was defined as $P < 0.05$.

Results

Baseline characteristics of patients at admission are shown in Table 1. Study population showed, as a whole, pulmonary hypertension and a mild increase of CO. Heart rate ($P=0.265$), systolic and mean systemic blood pressures (respectively $P=0.054$ and $P=0.134$), systolic PAP ($P=0.063$), mean PAP ($P=0.298$), cardiac output ($P=0.829$) and cardiac index ($P=0.851$) did not change significantly during the application of BiPAP through a face mask in the study population considered as a whole. After 30 min of NPPV application, no significant changes were observed in PaO₂ values when compared with PaO₂ following oxygen therapy alone. On the contrary, significant differences were found between pH, PaCO₂ and RR assessed during O₂ therapy alone and during NPPV (Table 2). Table 2 shows the results of echocardiograph evaluations (mean ± SD). Most of the 19 acute COPD patients did not have significant changes in Doppler findings, while six (31.5%) showed variations ≥ 15% of Doppler haemodynamic parameters during NPPV plus O₂ when compared with O₂ therapy alone. Four of these patients experienced

TABLE 2. Patients' measurements during O₂ supplementation and after 30 min of NPPV (mean \pm SD)

Parameters	O ₂ alone	O ₂ + NPPV	P value
pH	7.25 \pm 0.09	7.32 \pm 0.05	<0.001*
PaO ₂ (kPa)	8.71 \pm 1.11	8.75 \pm 1.01	0.870
PaCO ₂ (kPa)	10.25 \pm 1.76	8.57 \pm 1.09	<0.001*
RR (breaths min ⁻¹)	30 \pm 7	25 \pm 3	<0.001*
Mean BP (mmHg)	109.58 \pm 21.97	104.68 \pm 13.45	0.134
Systolic BP (mmHg)	150.25 \pm 31.27	141.50 \pm 19.74	0.064
HR (beats min ⁻¹)	100.42 \pm 16.09	97.84 \pm 16.35	0.265
PAMP (mmHg)	37.27 \pm 5.38	35.63 \pm 5.30	0.298
AcT/RVET	0.38 \pm 0.04	0.40 \pm 0.06	0.089
PASP (mmHg)	59.27 \pm 13.83	55.89 \pm 13.26	0.063
CO (l min ⁻¹)	6.59 \pm 1.93	6.65 \pm 2.03	0.829
CI (l min ⁻¹ m ⁻²)	3.53 \pm 0.86	3.56 \pm 0.97	0.851

*Statistically significant.

TABLE 3. Patients showing changes of $\geq 15\%$ in echocardiographic haemodynamic measurements after 30 min of NPPV administration plus O₂

Patient no.	PASP (mmHg)	PAMP (mmHg)	CI (l min ⁻¹ m ⁻²)	Systolic BP (mmHg)	Diastolic BP (mmHg)	HR (beats min ⁻¹)	RR (breaths min ⁻¹)	Mask tolerance	PaCO ₂ (kPa)	pH
3 O ₂	82	47.5	2.85	110	70	102	32		10.26	7.29
3 O ₂ + NPPV	82	45	2.25 (- 21%)	115	85	100	34	Poor	8.24	7.35
5 O ₂	NA	43	4.5	145	80	119	30		11.91	7.17
5 O ₂ + NPPV	NA	34 (- 21%)	3.5 (- 22%)	140	80	113	26	Good	7.84	7.33
6 O ₂	78	43	4.55	150	90	86	30		10.1	7.19
6 O ₂ + NPPV	66 (- 16%)	39	5.76 (+ 26%)	150	90	110	28	Good	8.67	7.29
10 O ₂	NA	32	4.3	150	80	99	48		13.73	7.15
10 O ₂ + NPPV	NA	25 (- 22%)	3.57 (- 17%)	160	90	91	36	Poor	10.46	7.25
11 O ₂	60.4	43	2.39	160	90	110	34		11.67	7.22
11 O ₂ + NPPV	61	43	3.2 (+ 33%)	140	80	82	26	Good	8.28	7.34
14 O ₂	45	NA	4.23	130	80	125	28		11.1	7.29
14 O ₂ + NPPV	37.3 (- 18%)	NA	3.63 (- 15%)	140	80	108	32	Poor	9.17	7.34

NA, not assessed.

consistent decreases in cardiac output, although cardiac output increased for the remaining two. Moreover, four patients showed a significant reduction of PASP and/or PAMP. Table 3 shows data for patients who had PAP and/or CI changes $\geq 15\%$ during NPPV. No correlation was found between decreased CO and baseline data. Conversely, three patients showing CO reduction had poor tolerance to mask ventilation and did not improve RR during NPPV. All the remaining patients showed good compliance with mask ventilation. All but two patients

(17/19) had a successful outcome with NPPV therapy (success rate 89.5%).

REPEATABILITY AND FEASIBILITY OF DOPPLER MEASUREMENTS

Both intraobserver and interobserver variability for PASP, PAMP and CI was less than 6%. One or more haemodynamic parameters were obtained by Doppler

echocardiography in most of the patients (16/18=89.5% of the study population). Tricuspid regurgitation was detected in 57%, though Doppler recording quality was adequate for velocity measurements in 47%. Pulmonary flow velocity profiles were obtained from 12 of the 19 COPD patients (63%). Also, tracings of these 12 patients sufficed for mPAP evaluation. Determination of cardiac output was possible in 89% of subjects.

Discussion

Initiation of invasive mechanical ventilation by endotracheal intubation may cause alteration of haemodynamics, especially in patients with AVF, because of acute exacerbation of COPD (6,36). In this clinical setting, a reduction in systemic venous return, the administration of sedative drugs before intubation and the effects of progressive hyperinflation–auto-PEEP and fluid depletion are considered to be the major contributors to hypotension (5,6,37,38). Recently, NPPV has been introduced as an alternative to endotracheal intubation in severe exacerbation of COPD (39,40). Although NPPV is widely used for patients with COPD and AVF, no mention of negative haemodynamic effects of NPPV appears in the English language literature (4,41).

In the present study, we show that, during the initiation of NPPV, some haemodynamic alteration may take place in individual COPD patients with AVF. In fact, four patients (21%) showed reduction of cardiac index during NPPV. It is worth noting that no patients had hypotension during NPPV administration. Indeed, NPPV application did not significantly change haemodynamics in most of our study population and, after 30 min of NPPV, the main haemodynamic data (cardiac output and cardiac index, pulmonary and systemic blood pressures) remained stable in 13 patients (68.4%). Moreover, arterial pH and $PaCO_2$ improved significantly during NPPV administration in all patients.

The maintenance of normal CO after 30 min of NPPV in the majority of the patients was not due to an increase in heart rate, as this remained stable or decreased. Also, in patients showing decreases of CO during NPPV, no correlation was found with baseline data. Nevertheless, three of these patients showed a poor tolerance to mask ventilation, as was, also confirmed by stable or increased RR. A lack of synchrony with ventilatory interface and consequent increased RR may have contributed to single patients developing increased hyperinflation and auto-PEEP (42). Furthermore, auto-PEEP is known to lead to decreased venous return and reduced CO (9). A change in stroke volume and/or peripheral resistance could be the mechanism behind the decrease in CI, but the lack of hypotension showed that in any case there were no important clinical consequences. Nevertheless, in AVF the CI is often elevated because of increased oxygen demand by the respiratory muscles and a drop in the CI value sometimes might represent an improvement, whereas an increase in CI might actually mean that the patient is working harder and needing more O_2 delivered (13,14). Since cardiopulmonary interactions during positive pressure ventilation are com-

plex, we cannot exclude the possibility that fluid depletion levels might have influenced CO reduction in some patients.

With regard to pulmonary circulation, at baseline pulmonary hypertension was observed in all the patients, but none of our 19 patients increased pulmonary pressures during NPPV, although in four cases a relevant reduction of PAP occurred.

To our knowledge, this study is the first that evaluates the haemodynamics of NPPV in COPD patients with AVF. Ambrosino *et al.* (43) reported haemodynamics data obtained through invasive methods performed on COPD patients in a clinically stable condition. Ambrosino *et al.* (43) found a significant fall in CO only when an external PEEP of 5 cmH₂O was administered. Nevertheless, patients with AVF due to severe exacerbation of COPD generally have higher auto-PEEP than those in a stable condition (44). Thus, the administration of equal levels of external PEEP may counterbalance or increase auto-PEEP if provided in acute or stable clinical settings (23,42).

We found that Doppler echocardiography is a well-tolerated, efficient evaluation method that can be used at NPPV initiation in acute COPD patients. It is well known that ultrasound examination of the heart is technically difficult in patients with hyperinflation of the lungs (45). However, expert and motivated echocardiograph technicians can obtain suitable images in up to 80–97% of patients (46,47). In our experience, the subcostal approach successfully detected adequate Doppler signals in a large proportion of COPD patients, thus confirming the results of Tramarin *et al.* (48).

In conclusion, our study suggests that the initiation of NPPV is haemodynamically safe for most patients with exacerbated COPD and AVF, although individual COPD patients with AVF may experience reduction in CI in spite of adequate levels of oxygen saturation during initiation of NPPV by facial mask. This event may be associated with patient inability to synchronize with mask ventilation. Consequently, we strongly suggest that particular caution should be used when applying NPPV with predetermined and fixed pressures since a decrease in CO may occur in patients who are hypotensive or have low fluid volume. In addition, to maximize mask ventilation tolerance, attempts to optimize EPAP and IPAP are preferable to scheduled ventilatory support settings. Finally, we recommended Doppler echocardiography as a useful tool for early detection of haemodynamic alterations due to NPPV application in patients with AVF.

Acknowledgement

The authors are grateful to Dr Roberto Tramarin, Fondazione S.Maugeri, Tradate, Italy for his suggestions and review of the manuscript.

References

1. Bott J, Carrol MP, Conway JH, Keilty SEJ, Ward EM, Brown AM, Paul EA *et al.* Randomized controlled trial of nasal ventilation in acute ventilatory failure due to

- chronic obstructive airways disease. *Lancet* 1993; **341**: 1555–1557.
2. Kramer N, Meyer TJ, Meharg J, Cece RD, Hill NS. Randomized prospective trial of noninvasive positive pressure ventilation in acute respiratory failure. *Am J Respir Crit Care Med* 1995; **151**: 1799–1806.
3. Brochard L, Mancebo J, Wysocki M, Lofaso F, Conti G, Rauss A, Simonneau G, Benito S, Gasparetto A, Lemaire F, Isabey D, Harf A. Noninvasive ventilation for acute exacerbations of chronic obstructive pulmonary disease. *N Engl J Med* 1995; **333**: 817–822.
4. Meduri GU. Noninvasive positive-pressure ventilation in patients with acute respiratory failure. *Clin Chest Med* 1996; **17**: 513–553.
5. Pinsky MR. Hemodynamic effects of artificial ventilation. In: Shoemaker WC, Ayres SM, Grenvik A, Holbrook PR, eds. *Textbook of Critical Care*. Philadelphia, PA: W. B. Saunders, 1995, pp. 911–923.
6. Keith RL, Pierson DJ. Complications of mechanical ventilation. *Clin Chest Med* 1996; **17**: 439–451.
7. Jardin F, Brun-Ney D, Hardy A, Aegerter P, Beauchet A, Bourdarias JP. Combined thermomodulation and two-dimensional echocardiographic evaluation of right ventricular function during respiratory support with PEEP. *Chest* 1991; **99**: 162–168.
8. Naeije R, Torbicki A. More on the noninvasive diagnosis of pulmonary hypertension: Doppler echocardiography revisited. *Eur Respir J* 1995; **8**: 1445–1449.
9. Ranieri VM, Dambrosio M, Brienza N. Intrinsic PEEP and cardiopulmonary interaction in patients with COPD and acute ventilatory failure. *Eur Respir J* 1996; **9**: 1283–1292.
10. Le Gall JR, Loirat P, Alperovitch A *et al.* A simplified acute physiology score for ICU patients. *Crit Care Med* 1984; **12**: 975–977.
11. Kelly BJ, Matthews MA. Prevalence and severity of neurological dysfunction in critically ill patients. Influence on need for continued mechanical ventilation. *Chest* 1993; **104**: 1818–1824.
12. MacNee W, Wathen CG, Flenley DC, Muir AL. The effects of controlled oxygen therapy on ventricular function in acute and chronic respiratory failure. *Am Rev Respir Dis* 1988; **137**: 1289–1295.
13. Degaute D, Domenighetti G, Naeije R, Vincent JL, Treyvaud D, Perret CI. Oxygen delivery in acute exacerbation of chronic obstructive pulmonary disease. Effects of controlled oxygen therapy. *Am Rev Respir Dis* 1981; **124**: 26–30.
14. Lejeune P, Mols P, Naeije R, Hallemans R, Melot C. Acute hemodynamic effects of controlled oxygen therapy in decompensated chronic obstructive pulmonary disease. *Crit Care Med* 1984; **12**: 1032–1035.
15. Gong H. Positive-pressure ventilation in the adult respiratory distress syndrome. *Clin Chest Med* 1982; **3**: 69–88.
16. Bone RC and the Coalition for Critical Care Excellence of the SCCM. Standards of evidence for the safety and effectiveness of critical care monitoring devices and related interventions. *Crit Care Med* 1995; **23**: 1756–1763.
17. Braghiroli A, Donner CF. Bi-level positive airway pressure. *Eur Respir Rev* 1992; **2**: 398–399.
18. Confalonieri M, Aiolfi S, Gandola L, Scartabellati A, Della Porta R, Parigi P. Severe exacerbations of chronic obstructive pulmonary disease treated with BiPAP by nasal mask. *Respiration* 1994; **61**: 310–316.
19. Confalonieri M, Parigi P, Scartabellati A, Aiolfi S, Scorsetti S, Nava S, Gandola L. Noninvasive mechanical ventilation improves the immediate and long-term outcome of COPD patients with acute respiratory failure. *Eur Respir J* 1996; **9**: 422–430.
20. Meduri GU, Turner RE, Abou-Shala N, Wunderink R, Tolley E. Noninvasive positive pressure ventilation via face mask. First-line intervention in patients with acute hypercapnic and hypoxemic respiratory failure. *Chest* 1996; **109**: 179–193.
21. Pennock BE, Kaplan PD, Carlin BW, Sabangan JS, Magovern JA. Pressure support ventilation with a simplified ventilatory support system administered with a nasal mask in patients with respiratory failure. *Chest* 1991; **100**: 1371–1376.
22. Vitacca M, Rubini F, Foglio K, Scalvini S, Nava S, Ambrosino N. Non-invasive modalities of positive pressure ventilation improve the outcome of acute exacerbations in COLD patients. *Intensive Care Med* 1993; **19**: 450–455.
23. Appendini L, Patessio A, Zanaboni S, Carbone M, Gukov B, Donner CF, Rossi A. Physiologic effects of positive end-expiratory pressure and mask pressure support during exacerbations of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1994; **149**: 1069–1076.
24. Himelman RB, Stulbarg M, Kircher B, Lee E, Kee L *et al.* Noninvasive evaluation of pulmonary artery pressure during exercise by saline-enhanced Doppler echocardiography in chronic pulmonary disease. *Circulation* 1989; **79**: 863–871.
25. Chan WL, Currie PJ, Seward JB, Hagler DJ, Mair DD, Tajik AJ. Comparison of three Doppler ultrasound methods in the prediction of pulmonary artery pressure. *J Am Coll Cardiol* 1987; **9**: 549–554.
26. Kitabakate A, Inoue M, Asao M, Masuyama T *et al.* Noninvasive evaluation of pulmonary hypertension by a pulsed Doppler technique. *Circulation* 1983; **68**: 302–309.
27. Mahan G, Dabestani A, Gardin J, Allfie A, Burn C, Henry W. Estimation of pulmonary artery pressure by Doppler echocardiography. *Circulation* 1984; **70** (Suppl II): II-363.
28. Hunstman LL, Stewart DK, Barnes SR *et al.* Non-invasive Doppler determination of cardiac output in man; clinical validation. *Circulation* 1983; **67**: 593–602.
29. Schena M, Clini E, Errera D, Quadri A. Echo-Doppler evaluation of left ventricular impairment in chronic cor pulmonale. *Chest* 1996; **109**: 1446–1451.
30. Spencer KT, Laing RM, Newman A, Barow KM, Shroff SG. Doppler and electromagnetic comparisons of instantaneous aortic flow characteristics in primates. *Circ Res* 1991; **68**: 1369–1377.

31. Jacobstein MD, Hirschfeld S, Winnie G *et al.* Ventricular interdependence in severe cystic fibrosis: A two-dimensional echocardiographic study. *Chest* 1981; **80**: 339–344.
32. Jezekova J, Michaljanic A, Jezek V. Natural variability of pulmonary hemodynamics. *Eur Heart J* 1987; **8** (Suppl II): II40.
33. Moulinier L, Venet T, Schiller N, Kurtz TW, Morris C Jr, Sebastian A. Measurement of aortic blood flow by Doppler echocardiography: day to day variability in normal subjects and applicability in clinical research. *J Am Coll Cardiol* 1991; **17**: 1326–1333.
34. Bland JM, Altman DG. Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* 1986; **i**: 307–310.
35. Pinsky MR. Defining the hypoxic threshold. *Crit Care Med* 1991; **19**: 147–149.
36. Miro AM, Pinsky MR. Heart–lung interaction. In: Tobin ed. *Textbook of Critical Care*. Philadelphia, PA: W. B. Saunders, 1995, pp. 911–923.
37. Pierson DJ. Complications associated with mechanical ventilation. *Crit Care Clin* 1990; **6**: 711–724.
38. Biondi JW, Schulman DS, Matthay RA. Effects of mechanical ventilation on right and left ventricular function. *Clin Chest Med* 1988; **9**: 55–71.
39. Elliott MW. Noninvasive ventilation in chronic obstructive pulmonary disease. *N Engl J Med* 1995; **333**: 870–871.
40. Ambrosino N. Noninvasive mechanical ventilation in acute respiratory failure. *Eur Respir J* 1996; **9**: 795–807.
41. Confalonieri M, Aiolfi S, Scartabellati A. Noninvasive ventilation. *Ann Intern Med* 1994; **121**: 984–985.
42. Rossi A, Polese G, Brandi G, Conti G. Intrinsic, positive end-expiratory pressure (PEEPi). *Intensive Care Med* 1995; **21**: 522–536.
43. Ambrosino N, Nava S, Torbicki A, Riccardi G, Fracchia C, Opasich C, Rampulla C. Haemodynamic effects of pressure support and PEEP ventilation by nasal route in patients with stable chronic obstructive pulmonary disease. *Thorax* 1993; **48**: 523–528.
44. Smith TC, Marini JJ. Impact of PEEP on lung mechanics and work of breathing in severe airflow obstruction. *J Appl Physiol* 1988; **65**: 1488–1499.
45. Morpurgo M, Saviotti M, Dickelè M, Casazza F, Torbicki A, Weitzenblum E, Zielinski J. Echocardiographic aspects of pulmonary artery hypertension in chronic lung disease. *Bull Eur Physiopathol Respir* 1984; **20**: 251–255.
46. Tramarin R, Torbicki A, Marchandise B, Laaban JP, Morpurgo M. Doppler echocardiographic evaluation of pulmonary artery pressure in chronic obstructive pulmonary disease. A European multicenter study. *Eur Heart J* 1991; **12**: 103–111.
47. Danchin N, Cornette EA, Henriquez A, Godenir JP, Ethevenot G, Polu JM, Sadoul P. Two dimensional echocardiographic assessment of the right ventricle in patients with chronic obstructive lung disease. *Chest* 1987; **92**: 229–233.
48. Tramarin R, Saviotti M, Colombo E *et al.* Echocardiographic and Doppler evaluation of pulmonary artery pressure in chronic lung disease. In: Morpurgo M *et al.* *Pathophysiology and Treatment of Pulmonary Circulation*. London: Springer, 1988, pp. 47–68.